**Bill Rooney** To: Subject: by kandy korn Friday, November 06, 2009 7:06:54 AM Date: Dr. Roonev. should be different from eack other if hybridization occured and That is correct. The parents will each plant being tested needs to be run seperately. The all have identical genotypes so they do not need to be tested individually. Matt ---- Original Message -----From: "Bill Rooney" <wlr@tamu.edu> To: "Patricia Klein" <pklein@tamu.edu> Cc: "Matthew Scott' 'Bartek" <mbartek@neo.tamu.edu> Sent: Thursday, November 5, 2009 1:46:01 PM GMT -06:00 US/Canada Central Subject: RE: by Trish: I expect that you've got hybrid corn, and seed derived from the cross of . Each seed would be different if KK DNA is actually present (which is not all that likely give what we see in the greenhouse). Matt correct me if this is wrong. I expect if you can run five different plants of the that would suffice and prove our point either way. Make sense? If not, let me know. Regards, Bill Dr. William L. Rooney Professor, Sorghum Breeding and Genetics Chair, Plant Release Committee Texas A&M University College Station, Texas 77843-2474 979 845 2151 -----Original Message-----From: Patricia Klein [mailto:pklein@tamu.edu] Sent: Thursday, November 05, 2009 11:44 AM To: Bill Rooney Subject: by kandy korn Bill I am a bit confused on the K marker work that you asked Natalie to do. Matt dropped off seed of the following: Thus he gave us three envelopes. My question is was there only one

x KK that you wanted us to check or is he sending us from several crosses? Before I have Natalie do anything

Bartek, Matthew Scott

From:

cross of

bulked

I want to know what we have. She and Matt both seemed a bit confused and I wasn't there to hear the conversation.

Thanks Trish

Dr. Patricia Klein Associate Professor Institute for Plant Genomics and Biotechnology TAMU 2123 Texas AgriLIFE Research Texas A&M University College Station, TX 77843-2123

phone: 979-862-6308 fax: 979-862-4790